# Post-Operative Instructions for Periodontal Surgery

## Work/ Exercise

Please go home and rest today. You may return to work tomorrow. If you choose to lie down, keep your head elevated on two pillows. Try to limit speaking for the first hour after surgery. Strenuous exercise should be avoided for 5 to 7 days.

## Medications

Take your antibiotics as prescribed. It is usually best <u>not</u> to take any medication on an empty stomach. You may take your prescription pain medication as needed, however, most patients find that **ibuprofen 600mg**. (Advil, Motrin) is very effective. It is helpful to take the first dose of ibuprofen immediately after surgery and before the anesthesia wears off. You may take a second dose as early as four hours after the first.

### **Ice Packs**

It is advantageous to place an ice pack on the cheek adjacent to the surgery site in a 5 minutes on/5 minutes off manner immediately after surgery. No ice should be used after the first 24 hours.

## **Periodontal Dressing**

A periodontal dressing is often placed around the treated teeth. Occasionally this dressing may become loose and fall off. If this occurs, simply throw the dressing away as this is not a problem and of no concern.

#### Food

Limit your diet today to liquids and very soft foods. Do not drink through a straw as this can cause bleeding. Cold beverage are advised for the first two or three hours. Soup or warm beverages may be consumed if only warm and not hot. Tomorrow you may return to a more normal soft diet being careful to chew only on the non-treated side.

## Brushing

Do not brush the <u>treated area</u> for one week following surgery. The other non-treated areas you may brush normally. On the second day after surgery, begin gently cleaning the treated teeth with a cotton swab dipped in Listerine. You may gently swab the teeth and gum-line or dressing.

## Rinsing

Do not rinse or spit today. Tomorrow begin rinsing two or three times a day with Listerine.

## Contacting Dr. Burns

In case you have any concerns after your procedure, you may contact Dr. Burns by calling our office number (404-929-9957). After hours and on weekends, **follow the voice mail prompt to activate our paging system.**